



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

The Construction of Monopoles

Citation for published version:

Braden, H & Enolskiy, V 2018, 'The Construction of Monopoles', *Communications in Mathematical Physics*, vol. 362, no. 2, pp. 547-570. <https://doi.org/10.1007/s00220-018-3199-4>

Digital Object Identifier (DOI):

[10.1007/s00220-018-3199-4](https://doi.org/10.1007/s00220-018-3199-4)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Communications in Mathematical Physics

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



THE CONSTRUCTION OF MONOPOLES

H.W. BRADEN AND V.Z. ENOLSKI

ABSTRACT. We show that the Higgs and gauge fields for a BPS monopole may be constructed directly from the spectral curve without having to solve the gauge constraint needed to obtain the Nahm data. The result is the analogue of the instanton result: given ADHM data one can reconstruct the gauge fields algebraically together with differentiation. Here, given the spectral curve, one can similarly reconstruct the Higgs and gauge fields. This answers a problem that has remained open since the discovery of monopoles.

1. INTRODUCTION

Despite the study of BPS monopoles being a mature subject, now over 35 years old, and having uncovered many remarkable results, a number of the original questions that sparked its development remain unanswered. They are hard. One can ask, for example, what the Higgs and gauge fields or their gauge invariant energy density are, but beyond the spherically symmetric coincident n -monopole solution and some partial results for $SU(2)$ charge 2 monopoles there are no explicit formulae. Although a number of general methods have been developed to address this, extending earlier analogous results for the construction of instantons (see below), these constructions typically involve a step (such as solving an ODE) that stymie attempts at an analytic solution. Numerical results based on these constructions and utilising the increase in computing power over the period has meant that we can understand a number of qualitative aspects of monopole behaviour; at the very least analytic solutions would give some control over these. In this paper we shall describe how to explicitly construct the Higgs and gauge fields for a monopole and circumvent those usually intractable steps; a number of new results will appear in the process.

We will focus here on $SU(2)$ charge- n monopoles. Our approach will assume from the outset that we know the spectral curve \mathcal{C} for the monopole. This curve appears in both Nahm's extension [13] of the ADHM construction of instantons and Ward's [16] use of the \mathcal{A}_k ansatz which Atiyah and Ward used in their instanton construction; it is also implicit in the Bäcklund transformation construction of Forgács, Horváth and Palla [8]. This curve gives a point in the moduli space of $SU(2)$ charge- n monopoles and we note that the relationship between this description of the moduli space and both Donaldson's rational map and Jarvis's rational map descriptions remains still poorly understood. In this paper we will work within the Nahm construction [9, 14]. This involves two potentially difficult steps: first, the solution of a first order (matrix) differential equation $\Delta^\dagger v = 0$; and second, the integration of appropriate bilinears of these solutions to give the Higgs and gauge fields. Now the operator Δ^\dagger is in turn constructed from Nahm data, matrices $T_i(s)$ ($i = 1, 2, 3$) that satisfy Nahm's equations $\dot{T}_1 = [T_2, T_3]$ (and cyclic) and certain boundary conditions that will be described in more detail below. Unfortunately Nahm data is hard to construct. Some years ago Ercolani and Sinha [7] showed how, using integrable systems techniques, one could solve for a gauge transform of the Nahm data directly from \mathcal{C} ; this theory has been further extended by the authors [2, 5]. Determining the gauge transformation is equivalent

to solving a further ODE and although its solution exists we don't know how to do this explicitly; we shall relate this to Donaldson's treatment [6] of the complex and real moment map description of the Nahm equations. Putting this difficulty to one side for the moment we show how integrable systems techniques and a *lesser known ansatz of Nahm* may be used to solve $\Delta^\dagger v = 0$, again up to the same gauge transformation. Not all of the solutions obtained are normalisable and in due course we show that constructing an appropriate projector to these is purely algebraic. But even with these solutions we must perform a number of integrations to obtain expressions for the sought after fields. An old work of Panagopoulos [15] is appropriate here: after some correction and small extensions we show that not only can all integrations be performed but the as yet undetermined gauge transformation combines within the bilinears into a term that is determined by the spectral curve. Rather than getting overly involved in the mathematics of Riemann surfaces that any example will necessitate (see for example [1, 3]) this paper will give general formulae for the fields directly in terms of the curve and a well-studied algebro-geometric object, the Baker-Akhiezer function. We conclude with a limited example showing how our construction yields a known result in the charge 2 setting; in a sequel paper we shall present the general results for the fields of the charge 2 monopole.

2. THE ADHMN CONSTRUCTION

In this section we recall the salient features of the ADHMN construction of monopoles sufficient to introduce our notation and the major features of the construction.

The equations we wish to solve are

$$(2.1) \quad D_i \Phi = \frac{1}{2} \sum_{j,k=1}^3 \epsilon_{ijk} F_{jk}, \quad i = 1, 2, 3,$$

for the gauge group $SU(2)$. Here Φ is the Higgs field, $F_{ij} = \partial_i A_j - \partial_j A_i + [A_i, A_j]$ is the curvature of the (spatial) connection of the gauge field $A_i(\mathbf{x})$ and D_i the covariant derivative $D_i \Phi = \partial_i \Phi + [A_i, \Phi]$, $\mathbf{x} = (x_1, x_2, x_3) \in \mathbb{R}^3$. These equations may be viewed as a reduction of the self-dual Yang Mills equations to three dimensions under the assumption that all fields are independent of time. Upon identifying the A_4 -component of the gauge field with the Higgs field Φ the four-dimensional Yang-Mills Lagrangian yields upon reduction the three dimensional Yang-Mills-Higgs Lagrangian

$$L = -\frac{1}{2} \text{Tr } F_{ij} F^{ij} + \text{Tr } D_i \Phi D^i \Phi.$$

We are interested in configurations minimizing the energy of the system. These are given by the *Bogomolny equation* (2.1) A solution with the boundary conditions

$$\left| \sqrt{-\frac{1}{2} \text{Tr } \Phi(r)^2} \right|_{r \rightarrow \infty} \sim 1 - \frac{n}{2r} + O(r^{-2}), \quad r = \sqrt{x_1^2 + x_2^2 + x_3^2},$$

is called a *monopole* of charge n .

Modifying the Atiyah-Drinfeld-Hitchin-Manin (ADHM) construction of instanton solutions to the (Euclidean) self-dual Yang-Mills equations Nahm introduced the operator

$$(2.2) \quad \Delta = \imath \frac{d}{dz} + x_4 - \imath T_4 + \sum_{j=1}^3 \sigma_j \otimes (T_j + \imath x_j 1_n),$$

where the $T_j(z)$ are $n \times n$ matrices and σ_j the Pauli matrices. Following the instanton construction the operator $\Delta^\dagger \Delta$ must commute with quaternions which happens if and only if $T_i^\dagger = -T_i$, $T_4^\dagger = -T_4$ and

$$(2.3) \quad \dot{T}_i = [T_4, T_i] + \frac{1}{2} \sum_{j,k=1}^3 \epsilon_{ijk} [T_j(s), T_k(s)].$$

Equations (2.3) are known as Nahm's equations; one often encounters them in the more familiar gauge with $T_4 = 0$. When $\Delta^\dagger \Delta$ commutes¹ with quaternions it is a positive operator; in particular this means that $(\Delta^\dagger \Delta)(z)$ is an invertible operator and consequently Δ has no zero modes. To describe monopoles the matrices $T_j(z)$ are further required to be regular for $z \in (-1, 1)$ and have simple poles at $z = \pm 1$, the residues of which define an irreducible n -dimensional representation of the $su(2)$ algebra. Hitchin's analysis [9][§2] of the equation $\Delta^\dagger \mathbf{v} = 0$ tells us that has two normalizable solutions and it is in terms of these that the Atiyah-Drinfeld-Hitchin-Manin-Nahm (ADHMN) construction gives the gauge and Higgs field solutions.

Theorem 2.1 (ADHMN). *The charge n monopole solution of the Bogomolny equation (2.1) is given by*

$$(2.4) \quad \Phi_{ab}(\mathbf{x}) = \imath \int_{-1}^1 dz z \mathbf{v}_a^\dagger(\mathbf{x}, z) \mathbf{v}_b(\mathbf{x}, z), \quad a, b = 1, 2,$$

$$(2.5) \quad A_{i ab}(\mathbf{x}) = \int_{-1}^1 dz \mathbf{v}_a^\dagger(\mathbf{x}, z) \frac{\partial}{\partial x_i} \mathbf{v}_b(\mathbf{x}, z), \quad i = 1, 2, 3, \quad a, b = 1, 2.$$

Here the two ($a = 1, 2$) $2n$ -column vectors $\mathbf{v}_a(\mathbf{x}, z) = (v_1^{(a)}(\mathbf{x}, z), \dots, v_{2n}^{(a)}(\mathbf{x}, z))^T$ form an orthonormal basis on the interval $z \in [-1, 1]$

$$(2.6) \quad \int_{-1}^1 dz \mathbf{v}_a^\dagger(\mathbf{x}, z) \mathbf{v}_b(\mathbf{x}, z) = \delta_{ab},$$

for the normalizable solutions to the Weyl equation

$$(2.7) \quad \Delta^\dagger \mathbf{v} = 0,$$

where

$$(2.8) \quad \Delta^\dagger = \imath \frac{d}{dz} + x_4 - \imath T_4 - \sum_{j=1}^3 \sigma_j \otimes (T_j + \imath x_j 1_n).$$

The normalizable solutions form a two-dimensional subspace of the full $2n$ -dimensional solution space to the formal adjoint equation (2.7). The $n \times n$ -matrices $T_j(z)$, $T_4(z)$, called Nahm data, satisfy Nahm's equation (2.3) and the $T_j(z)$ are required to be regular for $z \in (-1, 1)$ and have simple poles at $z = \pm 1$, the residues of which define an irreducible n -dimensional representation of the $su(2)$ algebra; further

$$(2.9) \quad T_i(z) = -T_i^\dagger(z), \quad T_4(z) = -T_4^\dagger(z), \quad T_i(z) = T_i^T(-z), \quad T_4(z) = T_4^T(-z).$$

Our strategy will be to solve (2.7) and determine those solutions that are normalizable. In what follows we shall denote by $V = (\mathbf{v}_1, \dots, \mathbf{v}_{2n})$ and similarly $W = (\mathbf{w}_1, \dots, \mathbf{w}_{2n})$ the $2n \times 2n$ fundamental matrices of solutions to $\Delta^\dagger \mathbf{v} = 0$ and $\Delta \mathbf{w} = 0$ respectively. Then V can be chosen to be $(W^\dagger)^{-1}$. As we have already remarked these are not all normalizable.

¹Throughout the superscript \dagger means conjugated and transposed. We will at times emphasise the vectorial nature of an object by printing this in bold, e.g for vector $\mathbf{a}^\dagger = \overline{\mathbf{a}}^T$.

According to the ADMHN theorem the Nahm data $T_j(z)$ expanded in the vicinity of the end point $z = 1 - \xi$ behaves as

$$T_j(1 - \xi) = -\imath \frac{l_j}{\xi} + O(1), \quad j = 1, 2, 3,$$

where (the Hermitian) l_j define the irreducible n -dimensional representation of the $su(2)$ Lie algebra, $[l_j, l_k] = \imath \epsilon_{jkl} l_l$. Then (2.7) behaves in the vicinity of the pole as

$$(2.10) \quad \left[\frac{d}{d\xi} - \frac{\sum_{j=1}^3 \sigma_j \otimes l_j}{\xi} \right] \mathbf{v}(\mathbf{x}, 1 - \xi) = 0.$$

One can show (see for example [17]) that $\sum_{j=1}^3 \sigma_j \otimes l_j$ has only two distinct eigenvalues, $\lambda_a = (n - 1)/2$ with multiplicity $n + 1$ and $\lambda_b = -(n + 1)/2$ with multiplicity $n - 1$. If \mathbf{a}_i are eigenvectors associated with λ_a ($i = 1, \dots, n + 1$), and \mathbf{b}_j eigenvectors associated with λ_b ($j = 1, \dots, n - 1$), then (2.10) has solutions $\xi^{\lambda_a} \mathbf{a}_i$ and $\xi^{\lambda_b} \mathbf{b}_j$. Therefore normalizable solutions must lie in the subspace with positive $\lambda_a = (n - 1)/2$ and so we require that $\mathbf{v}(\mathbf{x}, 1)$ is orthogonal to the subspace with eigenvalue $-(n + 1)/2$, i.e.

$$\lim_{z \rightarrow 1^-} \mathbf{v}(\mathbf{x}, z)^T \cdot \mathbf{b}_j = 0, \quad j = 1, \dots, n - 1.$$

These $n - 1$ conditions coming from the behaviour at $z = 1$ thus yield a $n + 1$ dimensional space of solutions to $\Delta^\dagger \mathbf{v} = 0$. A similar analysis at $z = -1$ again yields a further $n - 1$ constraints resulting in two normalisable solutions on the interval. We define a projector μ onto this subspace; this is the $2n \times 2$ matrix such that

$$V\mu = (\mathbf{v}_1, \mathbf{v}_2)$$

where \mathbf{v}_a ($a = 1, 2$) are the normalizable solutions (2.6). That is

$$\int_{-1}^1 dz \mu^\dagger V^\dagger V \mu = \mu^\dagger \left(\int_{-1}^1 dz V^\dagger V \right) \mu = 1_2.$$

Although $V = V(\mathbf{x}, z)$ and $\mathbf{v}_a = \mathbf{v}_a(\mathbf{x}, z)$ are both z -dependent we note that the matrix μ does not depend on z for we have from

$$0 = \Delta^\dagger(\mathbf{v}_1, \mathbf{v}_2) = \Delta^\dagger(V\mu) = (\Delta^\dagger V)\mu + \imath V \frac{d}{dz} \mu = \imath V \frac{d}{dz} \mu$$

and the (generic) invertibility of $V(\mathbf{x}, z)$ that $\dot{\mu} = 0$ and hence that $\mu = \mu(\mathbf{x})$. Thus to reconstruct the gauge and Higgs fields we must construct the projector μ that extracts from V the two normalizable solutions.

At this stage the integrations in (2.1) look intractable but work of Panagopoulos enables their evaluation. Define the Hermitian matrices

$$(2.11) \quad \mathcal{H} = -\sum_{j=1}^3 x_j \sigma_j \otimes 1_n, \quad \mathcal{F} = \imath \sum_{j=1}^3 \sigma_j \otimes T_j, \quad \mathcal{Q} = \frac{1}{r^2} \mathcal{H} \mathcal{F} \mathcal{H} - \mathcal{F}.$$

Then

Proposition 2.2 (Panagopoulos [15]).

$$(2.12) \quad \int dz \mathbf{v}_a^\dagger \mathbf{v}_b = \mathbf{v}_a^\dagger \mathcal{Q}^{-1} \mathbf{v}_b.$$

$$(2.13) \quad \int dz z \mathbf{v}_a^\dagger \mathbf{v}_b = \mathbf{v}_a^\dagger \mathcal{Q}^{-1} \left(z + \mathcal{H} \frac{x_i}{r^2} \frac{\partial}{\partial x_i} \right) \mathbf{v}_b.$$

$$(2.14) \quad \int \mathbf{v}_a^\dagger \frac{\partial}{\partial x_i} \mathbf{v}_b dz = \mathbf{v}_a^\dagger \mathcal{Q}^{-1} \left[\frac{\partial}{\partial x_i} + \mathcal{H} \frac{z}{r^2} x_i + \mathcal{H} \frac{\imath}{r^2} (\mathbf{x} \times \nabla)_i \right] \mathbf{v}_b.$$

These are proven in Appendix A. Interestingly, consideration of gauge invariance leads to new results that will be particularly useful in our later development. Recall that a gauge transformation acts on the normalizable solutions $(\mathbf{v}_1, \mathbf{v}_2) = V\mu$ from the right by $h(\mathbf{x}) \in SU(2)$. Then using (2.13)

$$(2.15) \quad \Phi = i \int_{-1}^1 dz z \mathbf{v}^\dagger \mathbf{v} = i \mu^\dagger \left[\int_{-1}^1 dz z V^\dagger V \right] \mu = i \left[\mu^\dagger V^\dagger (\mathcal{Q}^{-1} \left(z + \mathcal{H} \frac{x_i}{r^2} \frac{\partial}{\partial x_i} \right)) V \mu \right]_{z=-1}^{z=1}$$

and so a gauge transformation yields

$$\begin{aligned} h^{-1} \Phi h &= h^{-1} \left(i \int_{-1}^1 dz z \mathbf{v}^\dagger \mathbf{v} \right) h = h^{-1} \left(i \mu^\dagger \left[\int_{-1}^1 dz z V^\dagger V \right] \mu \right) h \\ &= h^{-1} \left(i \left[\mu^\dagger V^\dagger (\mathcal{Q}^{-1} \left(z + \mathcal{H} \frac{x_i}{r^2} \frac{\partial}{\partial x_i} \right)) V \mu \right]_{z=-1}^{z=1} \right) h \\ &= h^{-1} \Phi h + h^{-1} \left(i \left[\mu^\dagger V^\dagger \mathcal{Q}^{-1} \mathcal{H} V \mu \right]_{z=-1}^{z=1} \right) \frac{x_i}{r^2} \frac{\partial}{\partial x_i} h \end{aligned}$$

thus we must have

$$(2.16) \quad 0 = \left[\mu^\dagger V^\dagger \mathcal{Q}^{-1} \mathcal{H} V \mu \right]_{z=-1}^{z=1}.$$

Further, the transformation of the gauge field

$$(2.17) \quad A_i = \int_{-1}^1 dz \mathbf{v}^\dagger \frac{\partial}{\partial x_i} \mathbf{v} = \mu^\dagger V^\dagger \left(\mathcal{Q}^{-1} \left[\frac{\partial}{\partial x_i} + \mathcal{H} \frac{z}{r^2} x_i + \mathcal{H} \frac{\imath}{r^2} (\mathbf{x} \times \nabla)_i \right] \right) V \mu \Big|_{z=-1}^{z=1}$$

under a gauge transformation necessitates that we have

$$\begin{aligned} h^{-1} A_i h + h^{-1} \partial_i h &= h^{-1} \left(\int_{-1}^1 dz \mathbf{v}^\dagger \frac{\partial}{\partial x_i} \mathbf{v} \right) h = h^{-1} A_i h + h^{-1} \left(\mu^\dagger V^\dagger \mathcal{Q}^{-1} V \mu \right) \Big|_{z=-1}^{z=1} \partial_i h \\ &\quad + h^{-1} \left(\mu^\dagger V^\dagger \mathcal{Q}^{-1} \mathcal{H} V \mu \right) \Big|_{z=-1}^{z=1} \frac{\imath}{r^2} (\mathbf{x} \times \nabla)_i h. \end{aligned}$$

This will follow again as a result of (2.16) and the requirement of the ADHMN theorem 2.1 that $V\mu$ is normalised by

$$1_2 = \mu^\dagger \left(\int_{-1}^1 dz V^\dagger V \right) \mu = \mu^\dagger(\mathbf{x}) (V^\dagger(\mathbf{x}, z) \mathcal{Q}^{-1}(\mathbf{x}, z) V(\mathbf{x}, z)) \Big|_{z=-1}^{z=1} \mu(\mathbf{x}).$$

The z -independence of the projectors μ together with the fact that the poles of V lie only at the end points means that determining the projector is purely algebraic and reduces to the cancellation of poles.

Consideration of (2.16) leads to a new result which is the analogue of Hitchin's hermitian form introduced in his description of monopoles [9].

Theorem 2.3. *With the notation above,*

$$(2.18) \quad (V^\dagger \mathcal{Q}^{-1} \mathcal{H} V)(z) = \text{constant}, \quad (W^\dagger \mathcal{Q} \mathcal{H} W)(z) = \text{constant},$$

and consequently (2.16) holds true.

Proof. As these are essentially the inverses of each other we shall only prove the first. Using the notation and results of Appendix A we find

$$\begin{aligned}
\frac{d}{dz} (V^\dagger \mathcal{Q}^{-1} \mathcal{H} V) &= V^\dagger \left(\frac{d}{dz} [\mathcal{Q}^{-1} \mathcal{H}] - (\mathcal{H} + \mathcal{F}) \mathcal{Q}^{-1} \mathcal{H} - \mathcal{Q}^{-1} \mathcal{H} (\mathcal{H} + \mathcal{F}) \right) V \\
&= V^\dagger (\mathcal{D}(\mathcal{Q}^{-1}) \mathcal{H} + \mathcal{Q}^{-1} (\mathcal{H} + \mathcal{F}) \mathcal{H} - \mathcal{Q}^{-1} \mathcal{H} (\mathcal{H} + \mathcal{F})) V \\
&= V^\dagger (\mathcal{H} + \mathcal{Q}^{-1} [\mathcal{F}, \mathcal{H}]) V \\
&= 0
\end{aligned}$$

where we have made explicit use of (A.4) and (A.2). \square

Let us summarise what we have thus far. Given

- Nahm Data (needed to construct Δ^\dagger and \mathcal{Q}),
- the fundamental matrices V or W to $\Delta^\dagger \mathbf{v} = 0$ or $\Delta \mathbf{w} = 0$, where $V = (W^\dagger)^{-1}$,

we can algebraically solve for the projector μ and consequently obtain the Higgs (2.4) and gauge fields (2.17). Although this is the strategy of our solution we shall ultimately show that we only require this data up to a gauge transformation and that these are determined by the curve.

3. INTEGRABILITY AND A LESSER KNOWN ANSATZ OF NAHM

In this section we shall express both the Nahm data and fundamental matrices V and W in terms of the curve, clearly identifying what may be done explicitly and what is implicit. The key ingredient is integrability which has not been visible so far. We first make some general remarks, then turn to the construction of the Nahm data and fundamental matrices.

Upon setting (with $T_i^\dagger = -T_i$, $T_4^\dagger = -T_4$)

$$\alpha = T_4 + \imath T_3, \quad \beta = T_1 + \imath T_2, \quad L = L(\zeta) := \beta - (\alpha + \alpha^\dagger) \zeta - \beta^\dagger \zeta^2, \quad M = M(\zeta) := -\alpha - \beta^\dagger \zeta,$$

one finds

$$\begin{aligned}
\dot{T}_i &= [T_4, T_i] + \frac{1}{2} \sum_{j,k=1}^3 \epsilon_{ijk} [T_j(z), T_k(z)] \iff \dot{L} = [L, M] \\
(3.1) \quad &\iff \begin{cases} \left[\frac{d}{dz} - \alpha, \beta \right] = 0, \\ \frac{d(\alpha + \alpha^\dagger)}{dz} = [\alpha, \alpha^\dagger] + [\beta, \beta^\dagger]. \end{cases}
\end{aligned}$$

The first equivalence here is that of a Lax pair, suggesting an underlying integrable system, while the second equivalence expresses Nahm's equations in the form of a complex and a real equation (respectively) [6]. The complex Nahm equation is readily solved,

$$(3.2) \quad \beta g = g \nu, \quad \left(\frac{d}{dz} - \alpha \right) g = 0 \iff \beta = g \nu g^{-1}, \quad \alpha = \dot{g} g^{-1},$$

where ν is constant and generically diagonal, $\nu = \text{Diag}(\nu_1, \dots, \nu_n)$; by conjugating² by the constant matrix $g(0)$ we may assume $\beta(0) = \nu$ and $g(0) = 1_n$. It is the real equation that is more difficult. Define

$$(3.3) \quad h = g^\dagger g$$

² $\tilde{\beta} = g(0)^{-1} \beta g(0)$, $\tilde{g}(z) = g(0)^{-1} g(z)$, $\tilde{\alpha} = g(0)^{-1} \alpha g(0)$.

then

$$(3.4) \quad \dot{h}h^{-1} = g^\dagger(\alpha + \alpha^\dagger)g^{\dagger-1}, \quad h(0) = 1_n,$$

and the real equation yields the (possibly) nonabelian Toda equation

$$(3.5) \quad \frac{d}{dz}(\dot{h}h^{-1}) = [h\nu h^{-1}, \nu^\dagger].$$

In the monopole context Donaldson [6] proved the existence of a solution for the real equation with Nahm data. In terms of the Lax pair we are wishing to solve

$$(3.6) \quad \begin{aligned} (L - \eta)U &= 0, \\ \left[\frac{d}{dz} + M \right] U &= 0. \end{aligned}$$

The characteristic equation $P(\eta, \zeta) := \det(\eta - L(\zeta)) = 0$ defines our spectral curve \mathcal{C} which takes the form

$$(3.7) \quad P(\zeta, \eta) := \eta^n + a_1(\zeta)\eta^{n-1} + \dots + a_n(\zeta) = 0, \quad \deg a_k(\zeta) \leq 2k.$$

For large ζ we see that $\det(\eta/\zeta^2 - L/\zeta^2) \sim \prod_{i=1}^n (\eta/\zeta^2 + \nu_i^\dagger)$ and so $\eta/\zeta \sim -\nu_i^\dagger \zeta$. The curve \mathcal{C} is an n -sheeted cover of \mathbb{P}^1 of genus $g_{\mathcal{C}} = (n-1)^2$; we shall denote by $\{\infty_i\}_{i=1}^n$ the preimages of $\zeta = \infty$. Setting $U = g^{\dagger-1}\Phi$ we use the complex equation to transform (3.6) into a standard scattering equation for Φ ,

$$(3.8) \quad \left[\frac{d}{dz} - g^\dagger(\alpha + \alpha^\dagger)g^{\dagger-1} \right] \Phi = \zeta\nu^\dagger\Phi.$$

“Standard” here simply means that the matrix $\zeta\nu^\dagger$ on the right-hand side is z -independent. In terms of h we have (3.4) and

$$(3.9) \quad g^\dagger L g^{\dagger-1} = h\nu h^{-1} - \dot{h}h^{-1}\zeta - \nu^\dagger\zeta^2.$$

The point to note is that we can solve the standard scattering equation (3.8) explicitly in terms of the function theory of \mathcal{C} by what is known as a Baker-Akhiezer function [12], and so too $\dot{h}h^{-1}$ and (the gauge transform) $g^\dagger L g^{\dagger-1} = \widehat{\Phi} \text{Diag}(\eta_1, \dots, \eta_n) \widehat{\Phi}^{-1}$, where $\widehat{\Phi} := \widehat{\Phi}(z, P)$ ($P \in \mathcal{C}$) is the fundamental matrix of solutions to (3.8). Asymptotically the i -th column of $\widehat{\Phi}$ behaves as $\exp(z\zeta\nu_i^\dagger)$ and the Baker-Akhiezer function is defined by

$$(3.10) \quad \lim_{P=P(\zeta, \eta) \rightarrow \infty_i} \widehat{\Phi}(z, P) \exp(-z\zeta\nu_j^\dagger) = \text{Diag}(\delta_{ij}).$$

In the monopole context Ercolani and Sinha were the first to construct the Baker-Akhiezer function and use this to study Nahm data [7]; this theory was (corrected and) extended in [5] with a needed generalization of the Abel-Jacobi map proved in [4]. In the notation above these works explicitly construct the Baker-Akhiezer solutions to (3.8) as well³ as $\dot{h}h^{-1}$. Thus the Nahm data is only determined up to the gauge transformation $g^{\dagger-1}$ which is to satisfy the differential equation (3.4) for g , where the left-hand side is to be viewed as specified. In the gauge where $T_4 = 0$, equivalently $\alpha = \alpha^\dagger$, this takes the form

$$2\dot{g}^\dagger g^{\dagger-1} = \dot{h}h^{-1}$$

which we cannot solve in general. (In the Hitchin system context this is the analogue of having solved the complex moment map equation $\bar{\partial}_A \phi = 0$ and then being faced with solving the real moment map equation $\mu(A) = F_A + [\phi, \phi^\dagger] = 0$.) In what follows we will not need

³In the notation of Ercolani and Sinha we have $g^{\dagger-1} = C$ and $\dot{h}h^{-1} = -Q_0(z)$.

the specific construction of these functions and will simply denote the solution to (3.8) by Φ_{BA} .

Although we can only solve for Nahm data up to a gauge transformation, what can we say about the solutions to either $\Delta^\dagger \mathbf{v} = 0$ or $\Delta \mathbf{w} = 0$? Nahm again made the seminal ansatz; with some small changes better suited for the connections to integrability we may encode this in terms of the following theorem (proven in Appendix B).

Theorem 3.1 (Nahm [14]). *Let $|s\rangle$ be an arbitrarily normalized spinor not in $\ker(1_2 + \hat{\mathbf{u}}(\mathbf{x}) \cdot \boldsymbol{\sigma})$, with $\hat{\mathbf{u}}(\mathbf{x})$ a unit vector independent of z and $\boldsymbol{\sigma} = (\sigma_1, \sigma_2, \sigma_3)$. Then*

$$(3.11) \quad \mathbf{w} := \mathbf{w}(\zeta) = (1_2 + \hat{\mathbf{u}}(\mathbf{x}) \cdot \boldsymbol{\sigma}) e^{-\imath z[(x_1 - \imath x_2)\zeta - \imath x_3 - x_4]} |s\rangle \otimes U(z)$$

satisfies $\Delta \mathbf{w} = 0$ if and only if

$$(3.12) \quad 0 = (L(\zeta) - \eta) U(z),$$

$$(3.13) \quad 0 = \left(\frac{d}{dz} + M(\zeta) \right) U(z),$$

where

$$(3.14) \quad \eta = (x_2 - \imath x_1) - 2x_3\zeta - (x_2 + \imath x_1)\zeta^2,$$

and $L(\zeta)$ and $M(\zeta)$, as above, satisfy the Lax equation $\dot{L} = [L, M]$.

The appendix also shows that the curve has a real structure, that is, it is invariant under the anti-holomorphic involution:

$$(3.15) \quad \mathfrak{J} : (\zeta, \eta) \rightarrow \left(-\frac{1}{\bar{\zeta}}, -\frac{\bar{\eta}}{\bar{\zeta}^2} \right).$$

Given our previous discussion our approach is natural: we may solve

$$(3.16) \quad U(z) = g^{\dagger -1} \Phi_{BA}$$

in terms of the earlier (and unknown) gauge transformation $g^{\dagger -1}$ and the Baker-Akhiezer function. Prior to this other workers had sought to explicitly perform the integrations. (This approach is reviewed in Appendix C.)

It remains to describe the fundamental matrix W . Given a spectral curve and a position \mathbf{x} we substitute the corresponding value of η (given by (3.14)) into (3.7). This is an equation of degree $2n$ in ζ which we shall refer to as the *Atiyah-Ward constraint*, this equation having appeared in their work. The $2n$ solutions ζ_j give us $2n$ points on the curve $P_j := (\zeta_j, \eta_j)$ ($j = 1, \dots, n$) where η_j again follows from \mathbf{x} and (3.14). These $2n$ points come in n pairs of points related by the antiholomorphic involution \mathfrak{J} . To each point we have the associated values $\hat{\mathbf{u}}(\zeta_j)$ and for each of these we solve for $U(z)$ yielding a $2n \times 1$ matrix $\mathbf{w}(P_j)$. Taking each of the $2n$ solutions we obtain a $2n \times 2n$ matrix of solutions W . There may be non-generic points for which $\zeta_i = \zeta_j$ at which we modify this discussion by taking a derivative $\mathbf{w}'(P_j)$; these non-generic points correspond to points of bitangency of the spectral curve and appear in Hurtubise's study of the asymptotic behaviour of the Higgs field [11].

Summarising the insights of this section: the Nahm data, the fundamental matrices V or W to $\Delta^\dagger \mathbf{v} = 0$ or $\Delta \mathbf{w} = 0$, where $V = (W^\dagger)^{-1}$, and the Lax matrices L, M can be solved for in terms of the Baker-Akhiezer function up to the unknown gauge transformation $g^{\dagger -1}$. In terms of this gauge transformation $h = g^\dagger g$ satisfies the nonabelian Toda equation (3.5).

4. CONSTRUCTING THE GAUGE AND HIGGS FIELDS

Having reviewed the general formalism and established the first new result Theorem 2.3 we shall now extend this. Our aim is to establish

Theorem 4.1. *Given a spectral curve \mathcal{C} one may construct the gauge and Higgs fields directly.*

First, the form of (3.16) and (3.11) tells us that we may write

$$W = 1_2 \otimes g^{\dagger-1} \widehat{W}, \quad V = 1_2 \otimes g \widehat{V}, \quad \widehat{V} = \widehat{W}^{\dagger-1},$$

where

$$\widehat{w} = (1_2 + \widehat{u}(x) \cdot \sigma) e^{-i z [(x_1 - i x_2) \zeta - i x_3 - x_4]}|_S > \otimes \Phi_{BA}.$$

Here $\widehat{W} = (\widehat{w}_1, \dots, \widehat{w}_{2n})$ and \widehat{V} are explicitly expressible in terms of the spectral curve \mathcal{C} . Next observe that the operators appearing on the right-hand side of Proposition 2.2 take the form $\mathcal{Q}^{-1} \mathcal{O}$ and, because $g = g(z)$ is a function of z only, these indefinite integrals may be written as

$$\mu^{\dagger} V^{\dagger} \mathcal{Q}^{-1} \mathcal{O} V \mu = \mu^{\dagger} \widehat{V}^{\dagger} [(1_2 \otimes g^{\dagger}) \mathcal{Q}^{-1} (1_2 \otimes g)] \mathcal{O} \widehat{V} \mu = \mu^{\dagger} \widehat{V}^{\dagger} \mathcal{Q}'^{-1} (1_2 \otimes h) \mathcal{O} \widehat{V} \mu,$$

where we recall that $h = g^{\dagger} g$. Here, using the definitions (2.11),

$$\mathcal{Q}' = (1_2 \otimes g^{\dagger}) \mathcal{Q} (1_2 \otimes g^{\dagger-1}) := \frac{1}{r^2} \mathcal{H} \mathcal{F}' \mathcal{H} - \mathcal{F}', \quad \mathcal{F}' = i \sum_{j=1}^3 \sigma_j \otimes g^{\dagger} T_j g^{\dagger-1}.$$

Now using the definition of α, β we have⁴ from (3.9) that

$$\mathcal{F}' = \begin{pmatrix} \frac{1}{2} \dot{h} h^{-1} & -\nu^{\dagger} \\ i h \nu h^{-1} & -\frac{1}{2} \dot{h} h^{-1} \end{pmatrix},$$

and from our earlier remarks this may be reconstructed from the curve \mathcal{C} . Further \widehat{W} and \widehat{V} are determined by the Baker-Akhiezer function for the full range of $z \in [-1, 1]$, so allowing their expansions at the end-points and the evaluation of the integrals of Proposition 2.2. At this stage we see that the various impediments to reconstructing the gauge and Higgs fields, notably the need for the Nahm data to determine \mathcal{Q} and only being able to determine V up to a gauge transformation, have combined into the one unknown (matrix) $h(z)$. Certainly the Baker-Akhiezer function gives us $\dot{h} h^{-1}$ from which one can in principle solve the ODE for $h(z)$, but our aim is to reconstruct the gauge and Higgs fields without having to do this integration. We shall now show how we may reconstruct $h(z)$ and so prove the theorem.

5. DETERMINING $h(z)$

To proceed we must be a little more precise about our spectral curve, the nature of the coordinates (ζ, η) and the geometry of the Lax pair. In particular we need to specify the sections we are using in our construction. Hitchin's construction [9] shows that the spectral curve naturally lies in mini-twistor space: the spectral curve is an algebraic curve $\mathcal{C} \subset T\mathbb{P}^1$. If ζ is the inhomogeneous coordinate on the Riemann sphere then (ζ, η) are the standard local coordinates on $T\mathbb{P}^1$ defined by $(\zeta, \eta) \rightarrow \eta \frac{d}{d\zeta}$. The anti-holomorphic involution (3.15) endows $T\mathbb{P}^1$ with its standard real structure.

⁴ $\sum_j \sigma_j \otimes T_j = \frac{1}{2}(\sigma_1 + i\sigma_2) \otimes (T_1 - iT_2) + \frac{1}{2}(\sigma_1 - i\sigma_2) \otimes (T_1 + iT_2) + \sigma_3 \otimes T_3 = \sigma_- \otimes \beta + \sigma_3 \otimes T_3 - \sigma_+ \otimes \beta^{\dagger}.$

5.1. Bundle Structure and Sections. We wish to extend the definition of our Lax pair to yield a bundle on \mathbb{P}^1 . Let N and S denote the standard patches on \mathbb{P}^1 (here $0 \in N = \{\zeta \mid \zeta \neq \infty\}$) and denote by the same expressions the corresponding open sets for $T\mathbb{P}^1$. Take $L = L^N$, $M = M^N$, and we choose $(\zeta^S = 1/\zeta, \eta^S = \eta/\zeta^2)$

$$(5.1) \quad L^S(\zeta^S) = \frac{L^N(\zeta)}{\zeta^2}, \quad M^S(\zeta^S) = \alpha^\dagger - \frac{\beta}{\zeta} = M^N(\zeta) - \frac{L^N(\zeta)}{\zeta}.$$

We have the reality conditions

$$(5.2) \quad L^{S\dagger}(\zeta^S) = -L^N(-1/\bar{\zeta}), \quad M^{S\dagger}(\zeta^S) = -M^N(-1/\bar{\zeta}).$$

Away from the branch points of the spectral curve the eigenvectors⁵ U_j^N are linearly independent and can be organized into a maximal rank matrix $U^N = (U_1^N, U_2^N, \dots, U_n^N)$ so that $L^N = U^N D^N (U^N)^{-1}$, with diagonal $D^N = \text{Diag}(\eta_1^N, \eta_2^N, \dots, \eta_n^N)$. Viewing our curve \mathcal{C} as an n -fold branched cover of \mathbb{P}^1 the columns U_j^N correspond to the various sheets of the cover. Similarly, $L^S = U^S D^S (U^S)^{-1}$. From (5.1) we see that the corresponding eigenvectors are proportional: $U^S = U^N F$ with diagonal $F = \text{diag}(F_1, F_2, \dots, F_n)$. It also follows from (5.1) and $(d/dz + M^{N,S})U^{N,S} = 0$ that

$$(5.3) \quad \frac{d}{dz} F = \frac{D^N}{\zeta} F.$$

Thus F is diagonal with its diagonal elements of the form $F_i(z, \zeta, \eta) = f_i(\zeta) e^{z\eta_i/\zeta}$. For the case we consider the spectral curve has monodromy permuting all of its sheets (and thus all of its eigenvectors $U_j^{N,S}$); thus all functions the $f_i(\zeta)$ must be equal and so $F_i(z, \zeta, \eta) = f(\zeta) e^{z\eta_i/\zeta}$. Let us now focus on the I -th row of U^N : this defines a function Ψ_I^N on \mathcal{C} everywhere outside $\zeta = \infty$. Similarly, the I -th row of U^S defines a function Ψ_I^S on \mathcal{C} everywhere besides $\zeta = 0$. These two functions are related by $\Psi_I^S = \Psi_I^N f(\zeta) e^{z\eta/\zeta}$; in other words the pair of I -th rows of U^N and U^S define a section of the bundle over \mathcal{C} with transition function $f(\zeta) \exp(z\eta/\zeta)$. Now by the Birkhoff-Grothendieck theorem a line bundle over \mathbb{P}^1 must be of the form $\mathcal{O}(r)$ for some integer r , and so by a change of trivialization we can take $f(\zeta) = 1/\zeta^r$.

At this point we come to different choices in specifying the sections U . Let L^z be the line bundle with transition function $e^{z\eta/\zeta}$. For Hitchin $z \in [0, 2]$ with $z = 1$ the point where $L^1 \otimes \pi^* \mathcal{O}(n-1)$ has n sections with real structure. Hitchin's sections [9] are defined just in terms of the two patches N, S and $r = n-1$ in the above. The Baker-Akhiezer construction of [5, 7] describes the flow in terms of $s \in [-1, 1]$ ($z = s+1$) and a line bundle \mathcal{L}_δ corresponding to a nonspecial divisor $\delta = \sum_{i=1}^{g_C+n-1} \delta_i$ of degree $g_C + n - 1 = \deg \pi^* \mathcal{O}(n-1)$. The line bundles \mathcal{L}_δ and $L^1 \otimes \pi^* \mathcal{O}(n-1)$ are linearly equivalent. On $\mathcal{C} \setminus \{\infty_i\}_{i=1}^n$ the Baker-Akhiezer function is meromorphic with poles in δ ; the transition functions for this line bundle are around each of the δ_i . Thus in the NS transition function we have $r = 0$. In what follows we set

$$(5.4) \quad F = \text{Diag} \left(e^{z\eta_i/\zeta} / \zeta^l \right)$$

where $l = n-1$ when describing Hitchin's choice of section and $l = 0$ when describing the Baker-Akhiezer sections.

⁵We will use capital Roman letters $\{I, J, \dots\}$ for rows and lower case Roman letters $\{i, j, \dots\}$ for columns.

5.2. Identifying $h(z)$. The bundle structure just described enables us to identify $h(z)$. First observe that if $[d/dz + M]U = 0$ then $[d/dz - M^\dagger]U^{\dagger-1} = 0$. Using this together with (5.2) shows $[d/dz + M^S(\zeta^S)]U^N(z, -1/\bar{\zeta})^{\dagger-1} = 0$ and consequently that

$$(5.5) \quad \mathcal{D}^S(\zeta^S) := U^N(z, -1/\bar{\zeta})^\dagger U^S(z, \zeta^S)$$

is a z -independent matrix. Now

$$\begin{aligned} \mathcal{D}^S(\zeta^S) D^S(\zeta^S) &= U^N(z, -1/\bar{\zeta})^\dagger L^S(\zeta^S) U^S(z, \zeta^S) = - (L^N(-1/\bar{\zeta}) U^N(z, -1/\bar{\zeta}))^\dagger U^S(z, \zeta^S) \\ &= -D^N(-1/\bar{\zeta})^\dagger \mathcal{D}^S(\zeta^S) = D^S(\zeta^S) \mathcal{D}^S(\zeta^S). \end{aligned}$$

Using the fact that the diagonal matrices $D^{N,S}$ have distinct entries for generic (ζ, η) we deduce that \mathcal{D}^S is diagonal. We similarly define the constant diagonal matrix

$$(5.6) \quad \mathcal{D}^N(\zeta) = (-1)^l U^S(z, -1/\bar{\zeta}^S)^\dagger U^N(z, \zeta).$$

Then from $U^S = U^N F$ on an overlap we have

$$\mathcal{D}^S(\zeta^S) = \left(U^S(z, -\frac{1}{\bar{\zeta}^S}) F(z, -\frac{1}{\bar{\zeta}})^{-1} \right)^\dagger U^N(z, \zeta) F(z, \zeta) = (-1)^l F(z, -\frac{1}{\bar{\zeta}})^{\dagger-1} \mathcal{D}^N(\zeta) F(z, \zeta).$$

Upon using that all of the matrices here are diagonal and that under the antiholomorphic involution (3.15) $\eta/\zeta \rightarrow \bar{\eta}/\bar{\zeta}$ we have

$$\mathcal{D}^S = (-1)^l f(\zeta)/f(-1/\bar{\zeta})^\dagger \mathcal{D}^N = \frac{1}{\zeta^{2l}} \mathcal{D}^N.$$

In the Hitchin setting $\{\mathcal{D}^N, \mathcal{D}^S\}$ yield a section of $\pi^* \mathcal{O}(2n-2)$; such a section takes the form $c_0 \eta^{n-1} + c_1(\zeta) \eta^{n-2} + \dots + c_{n-1}(\zeta)$ where $c_l(\zeta)$ is of degree $2l$ in ζ . In our setting we get a regular function constant in z .

5.3. Expressing $h(z)$ in terms of the Baker-Akhiezer function. Now with $U = U^N = g(z)^{\dagger-1} \hat{\Phi}(z, \zeta)$, where $\hat{\Phi}(z, \zeta)$ is the $n \times n$ matrix whose columns⁶ are the Baker-Akhiezer functions for the preimages of ζ , we have

$$\mathcal{D}^N(\zeta) = \left[U^N(z, -\frac{1}{\bar{\zeta}}) \text{Diag}(e^{z\bar{\eta}_i/\bar{\zeta}}) \right]^\dagger g^{\dagger-1} \hat{\Phi}(z, \zeta) = \text{Diag}(e^{z\eta_i/\zeta}) \hat{\Phi}(z, -\frac{1}{\bar{\zeta}})^\dagger h(z)^{-1} \hat{\Phi}(z, \zeta).$$

Now the left-handside is z -independent, so we may evaluate this at $z = 0$ using $h(0) = 1_n$ to give

$$\mathcal{D}^N(\zeta) = \hat{\Phi}(0, -\frac{1}{\bar{\zeta}})^\dagger \hat{\Phi}(0, \zeta).$$

Employing this we obtain

$$(5.7) \quad h(z) = \hat{\Phi}(z, \zeta) \text{Diag}(e^{z\eta_i/\zeta}) \hat{\Phi}(0, \zeta)^{-1} \hat{\Phi}(0, -\frac{1}{\bar{\zeta}})^\dagger \hat{\Phi}(z, -\frac{1}{\bar{\zeta}})^\dagger.$$

Again using the diagonality of \mathcal{D}^N this expression shows that $h(z)^\dagger$ is obtained by the interchange of $\zeta \rightarrow -\frac{1}{\bar{\zeta}}$; but $h(z)$ is independent of ζ and so we see that $h(z)$ is hermitian, as is required. Finally, using the independence of $h(z)$ on the value of ζ being used, upon taking ζ to infinity and using (3.10) and the hermiticity of h we may rewrite this to give

Theorem 5.1. *Let $\hat{\Phi}(z, 0)$ be the Baker-Akhiezer function with (I, i) -entry $\hat{\Phi}_I(z, 0_i)$, where $0_i = \mathcal{J}(\infty_i)$, then*

$$(5.8) \quad h(z) = \hat{\Phi}(z, 0) \hat{\Phi}(0, 0)^{-1}.$$

⁶ $\hat{\Phi}(z, \zeta) := (\Phi_1(z, P_1), \dots, \Phi_n(z, P_n))$, where $P_i = (\zeta, \eta_i)$.

We remark that the inverse of $\widehat{\Phi}(z, \zeta)$ may be constructed with the dual Baker-Akhiezer function; further, we see that this expression is independent of the ordering of sheets that is implicit in the Baker-Akhiezer function.

At this stage we have established Theorem 4.1 circumventing the actual construction of Nahm data. To determine the Nahm data one must solve for $h(z) = \widehat{\Phi}(z, 0)\widehat{\Phi}(0, 0)^{-1} = g^\dagger g$ where h and g must satisfy (3.4). This constraint makes finding g more difficult than simply a matrix factorization problem; although Cholesky factorization enables one to factorize h such a factorization is only defined up to a unitary transformation which is determined by (3.4). An interesting question is whether the function theory of \mathcal{C} might help in this determination.

6. EXAMPLE

We shall give an example of Theorem 5.1 for the case $n = 2$. The spectral curve for $n = 2$ was constructed by Hurtubise [10] and we shall employ the Ercolani-Sinha [7] form

$$(6.1) \quad 0 = \eta^2 + \frac{K^2}{4} (\zeta^4 + 2(k^2 - k'^2)\zeta^2 + 1),$$

where $K = K(k)$ is a complete elliptic integral, and η is related to the spatial coordinates by

$$(6.2) \quad \eta = (x_2 - ix_1) - 2\zeta x_3 - (x_2 + ix_1)\zeta^2.$$

It was known that the solutions

$$(6.3) \quad \begin{aligned} f_1(z) &= K \frac{\operatorname{dn} Kz}{\operatorname{cn} Kz} = \frac{\pi\theta_2\theta_3}{2} \frac{\theta_3(z/2)}{\theta_2(z/2)}, & f_2(z) &= Kk' \frac{\operatorname{sn} Kz}{\operatorname{cn} Kz} = \frac{\pi\theta_3\theta_4}{2} \frac{\theta_1(z/2)}{\theta_2(z/2)}, \\ f_3(z) &= Kk' \frac{1}{\operatorname{cn} Kz} = \frac{\pi\theta_2\theta_4}{2} \frac{\theta_4(z/2)}{\theta_2(z/2)} \end{aligned}$$

to the spinning top equations $\dot{f}_1 = f_2 f_3$ (and cyclic) gave solutions to the Nahm equations via $T_j(z) = \frac{\sigma_j}{2i} f_j(z)$ and the work of [7] and (with corrections in) [5] derived these from first principles. As we have noted the construction proceeds via the Baker-Akhiezer function but this only yields the Nahm data up to a gauge transformation $g^{\dagger-1}$. In the works cited the differential equation for this gauge transformation was solved explicitly and we will compare the expression for $h(z)$ given by Theorem 5.1 with that obtained from the solution of the differential equation. For higher charges the associated differential equation has not been solved.

The Baker-Akhiezer function for the problem at hand takes the form

$$(6.4) \quad \Phi(z, P) = \chi(P) \begin{pmatrix} -\theta_3(\alpha(P))\theta_2(\alpha(P) - z/2) \\ \theta_1(\alpha(P))\theta_4(\alpha(P) - z/2) \end{pmatrix} \frac{e^{\beta_1(P)z}}{\theta_2(z/2)}$$

where

$$(6.5) \quad \chi(P) = \frac{\theta_2(1/4)\theta_3(1/4)}{\theta_3(0)\theta_1(\alpha(P) - 1/4)\theta_4(\alpha(P) + 1/4)}, \quad \beta_1(P) = \frac{1}{4} \left\{ \frac{\theta'_1(\alpha(P))}{\theta_1(\alpha(P))} + \frac{\theta'_3(\alpha(P))}{\theta_3(\alpha(P))} \right\},$$

and we have the Abel map $\alpha(P) = \int_{\infty_1}^P d\zeta/4\eta$ for $P = (\zeta, \eta)$. Then with $P_{1,2}$ corresponding to $(\zeta, \pm\eta)$ we have

$$(6.6) \quad \widehat{\Phi}(z, \zeta) = \begin{pmatrix} -\theta_3(\alpha(P_1))\theta_2(\alpha(P_1) - z/2) & -\theta_3(\alpha(P_2))\theta_2(\alpha(P_2) - z/2) \\ \theta_1(\alpha(P_1))\theta_4(\alpha(P_1) - z/2) & \theta_1(\alpha(P_2))\theta_4(\alpha(P_2) - z/2) \end{pmatrix} \\ \times \begin{pmatrix} \frac{\chi(P_1)e^{\beta_1(P_1)z}}{\theta_2(z/2)} & 0 \\ 0 & \frac{\chi(P_2)e^{\beta_1(P_2)z}}{\theta_2(z/2)} \end{pmatrix}.$$

To evaluate this for $0_{1,2}$ we note (from [5]) that $\alpha(0_1) = -\tau/2$, $\alpha(0_2) = -1/2$. Using $\theta_4(\tau/2) = 0 = \theta_2(1/2)$ we may simplify (5.8) to

$$h(z) = \begin{pmatrix} \frac{\theta_2(\alpha(0_1) - z/2)}{\theta_2[0_1]} & -\frac{\theta_3[0_2]\theta_2(\alpha(0_2) - z/2)}{\theta_1[0_2]\theta_4[0_2]} \\ -\frac{\theta_1[0_1]\theta_4(\alpha(0_1) - z/2)}{\theta_3[0_1]\theta_2[0_1]} & \frac{\theta_4(\alpha(0_2) - z/2)}{\theta_4[0_2]} \end{pmatrix} \begin{pmatrix} \frac{\theta_2 e^{\beta_1(0_1)z}}{\theta_2(z/2)} & 0 \\ 0 & \frac{\theta_2 e^{\beta_1(0_2)z}}{\theta_2(z/2)} \end{pmatrix} \\ = \begin{pmatrix} \frac{\theta_2(\tau/2 + z/2)}{\theta_2(\tau/2)} & \frac{\theta_3(1/2)\theta_2(1/2 + z/2)}{\theta_1(1/2)\theta_4(1/2)} \\ \frac{\theta_1(\tau/2)\theta_4(\tau/2 + z/2)}{\theta_3(\tau/2)\theta_2(\tau/2)} & \frac{\theta_4(1/2 + z/2)}{\theta_4(1/2)} \end{pmatrix} \begin{pmatrix} \frac{\theta_2 e^{\beta_1(0_1)z}}{\theta_2(z/2)} & 0 \\ 0 & \frac{\theta_2 e^{\beta_1(0_2)z}}{\theta_2(z/2)} \end{pmatrix} \\ = \begin{pmatrix} \frac{B(z/2)\theta_3(z/2)}{B(0)\theta_3} & -\frac{\theta_4\theta_1(z/2)}{\theta_2\theta_3} \\ -\frac{B(z/2)\theta_4\theta_1(z/2)}{B(0)\theta_2\theta_3} & \frac{\theta_3(z/2)}{\theta_3} \end{pmatrix} \begin{pmatrix} \frac{\theta_2 e^{\beta_1(0_1)z}}{\theta_2(z/2)} & 0 \\ 0 & \frac{\theta_2 e^{\beta_1(0_2)z}}{\theta_2(z/2)} \end{pmatrix}$$

where $B(v) = \exp(-i\pi[v + \tau/4])$

$$= \frac{1}{\theta_3\theta_2(z/2)} \begin{pmatrix} \theta_2\theta_3(z/2) & -\theta_4\theta_1(z/2) \\ -\theta_4\theta_1(z/2) & \theta_2\theta_3(z/2) \end{pmatrix} \begin{pmatrix} e^{\beta_1(0_1)z - i\pi z/2} & 0 \\ 0 & e^{\beta_1(0_2)z} \end{pmatrix}.$$

Now upon using the quasi-periodicity of the theta functions,

$$\frac{\theta'_1(-\tau/2)}{\theta_1(-\tau/2)} = i\pi - \frac{\theta'_4}{\theta_4}, \quad \frac{\theta'_3(-\tau/2)}{\theta_3(-\tau/2)} = i\pi - \frac{\theta'_2}{\theta_2},$$

we find

$$\beta(0_1) = \frac{i\pi}{2} - \frac{1}{4} \left\{ \frac{\theta'_2}{\theta_2} + \frac{\theta'_4}{\theta_4} \right\} = \frac{i\pi}{2}, \quad \beta(0_2) = -\frac{1}{4} \left\{ \frac{\theta'_2}{\theta_2} + \frac{\theta'_4}{\theta_4} \right\} = 0.$$

Here we have used that $\theta_{2,4}$ are even and so $\theta'_{2,4} = 0$. Thus

$$h(z) = \frac{1}{\theta_3\theta_2(z/2)} \begin{pmatrix} \theta_2\theta_3(z/2) & -\theta_4\theta_1(z/2) \\ -\theta_4\theta_1(z/2) & \theta_2\theta_3(z/2) \end{pmatrix}.$$

In terms of (6.3) and using $K = \pi\theta_3^2/2$ we arrive at

$$(6.7) \quad h(z) = \frac{1}{K} \begin{pmatrix} f_1 & -f_2 \\ -f_2 & f_1 \end{pmatrix}, \quad h^{-1}(z) = \frac{1}{K} \begin{pmatrix} f_1 & f_2 \\ f_2 & f_1 \end{pmatrix}.$$

We wish to compare (6.7) with $h^{-1} = C^\dagger C$ where C is the solution to the differential equation

$$C^{-1}\dot{C} = \dot{C}C^{-1} = \frac{f_3}{2} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}.$$

Then

$$C = C^\dagger = C^T = \begin{pmatrix} F(z) & G(z) \\ G(z) & F(z) \end{pmatrix},$$

with solution

$$F = \cosh \left(\int_0^z f_3(s)ds/2 \right) = [p(z) + 1/p(z)]/2, \quad G = \sinh \left(\int_0^z f_3(s)ds/2 \right) = [p(z) - 1/p(z)]/2,$$

where⁷

$$p(z) = \exp \left(\int_0^z f_3(s) ds/2 \right) = \exp \left(k' K \int_0^z \frac{ds}{\operatorname{cn} Kz} \right) = \left[\frac{\operatorname{dn} Kz + k' \operatorname{sn} Kz}{\operatorname{cn} Kz} \right]^{1/2}.$$

Now

$$(6.8) \quad G^2(z) = \frac{1}{2} \left(\frac{\operatorname{dn}(Kz; k)}{\operatorname{cn}(Kz; k)} - 1 \right) = \frac{1}{2} \left(\frac{f_1}{K} - 1 \right), \quad 2F(z)G(z) = k' \frac{\operatorname{sn}(Kz; k)}{\operatorname{cn}(Kz; k)} = \frac{f_2}{K},$$

$$F^2 - G^2 = 1, \quad F^2 + G^2 = \frac{f_1}{K},$$

and consequently $C^2(z) = \frac{1}{K} \begin{pmatrix} f_1 & f_2 \\ f_2 & f_1 \end{pmatrix}$, so verifying (6.7).

ACKNOWLEDGEMENTS

HWB thanks Sergey Cherkis, Derek Harland and Paul Sutcliffe for helpful conversations.

APPENDIX A. THE PANAGOPOULOS FORMULAE

Panagopolous [15] introduced three formulae to evaluate integrals appearing in the ADHMN construction giving a proof for one of these; his other formulae were checked against a calculation on one axis. We shall prove and slightly modify here the Panagopolous formulae, extending his method to the case x_4, T_4 possibly nonzero and correcting an ordinary differential operator by a partial differential operator (which reduces to the former on an axis).

Panagopolous's approach to evaluate integrals

$$\int dz \, v_a^\dagger \mathcal{A} v_b$$

for any given operator \mathcal{A} and any two solutions $v_{a,b}$ of $\Delta^\dagger v = 0$ is to seek an operator \mathcal{B} such that

$$(A.1) \quad v_a^\dagger \mathcal{A} v_b = \frac{d}{dz} (v_a^\dagger \mathcal{B} v_b),$$

so giving a primitive. Using (2.7, 2.11) then if v is any solution of $\Delta^\dagger v = 0$ we have that

$$1_{2n} \frac{d}{dz} v = [\imath x_4 + T_4 - (\mathcal{H} + \mathcal{F})] v,$$

whence

$$v_a^\dagger \mathcal{A} v_b = \frac{d v_a^\dagger}{dz} \mathcal{B} v_b + v_a^\dagger \frac{d \mathcal{B}}{dz} v_b + v_a^\dagger \mathcal{B} \frac{d v_b}{dz} = v_a^\dagger \left(\frac{d \mathcal{B}}{dz} - [T_4, \mathcal{B}] - (\mathcal{H} + \mathcal{F}) \mathcal{B} - \mathcal{B} (\mathcal{H} + \mathcal{F}) \right) v_b.$$

The diagonal term with x_4 cancels here. Thus we seek to relate the operators \mathcal{A} and \mathcal{B} by

$$\mathcal{A} = \mathcal{D}(\mathcal{B}) := \frac{d \mathcal{B}}{dz} - [T_4, \mathcal{B}] - (\mathcal{H} + \mathcal{F}) \mathcal{B} - \mathcal{B} (\mathcal{H} + \mathcal{F}).$$

where we have defined the operator \mathcal{D} .

We will use the following relations (recall T_4 is shorthand for $1_2 \otimes T_4$),

$$(A.2) \quad \mathcal{Q} \mathcal{H} = \mathcal{H} \mathcal{F} - \mathcal{F} \mathcal{H} = [\mathcal{H}, \mathcal{F}] = -\mathcal{H} \mathcal{Q}, \quad \mathcal{H} = \mathcal{Q}^{-1} [\mathcal{H}, \mathcal{F}], \quad [\mathcal{H}, T_4] = 0,$$

$$x_i (\partial_i \mathcal{H}) = \mathcal{H}, \quad \mathcal{H}^2 = r^2 1_{2n},$$

⁷ Here we have made use of

$$\int \frac{du}{\operatorname{cn} u} = \frac{1}{k'} \ln \frac{\operatorname{dn} u + k' \operatorname{sn} u}{\operatorname{cn} u}.$$

$$\mathcal{F}^2 = -1_2 \otimes \sum_{i=1}^3 T_i T_i - \imath \sum_{i,j,k=1}^3 \epsilon_{ijk} \sigma_k \otimes T_i T_j,$$

$$\frac{d\mathcal{F}}{dz} - [T_4, \mathcal{F}] = \imath \sum_{i,j,k=1}^3 \epsilon_{ijk} \sigma_k \otimes T_i T_j.$$

Therefore

$$(A.3) \quad 0 = \left[\mathcal{F}^2 + \frac{d\mathcal{F}}{dz} - [T_4, \mathcal{F}], \mathcal{H} \right].$$

We now establish the integrals in (2.2).

Proposition A.1. *With $\mathcal{Q} = \frac{1}{r^2} \mathcal{H} \mathcal{F} \mathcal{H} - \mathcal{F}$ then*

$$(A.4) \quad \mathcal{D}(\mathcal{Q}^{-1}) = \frac{d\mathcal{Q}^{-1}}{dz} - [T_4, \mathcal{Q}^{-1}] - (\mathcal{H} + \mathcal{F}) \mathcal{Q}^{-1} - \mathcal{Q}^{-1} (\mathcal{H} + \mathcal{F}) = 1_{2n}$$

and we have the antiderivative (2.12)

$$\int dz \mathbf{v}_a^\dagger \mathbf{v}_b = \mathbf{v}_a^\dagger \mathcal{Q}^{-1} \mathbf{v}_b.$$

Proof. In this case $\mathcal{A} = 1_{2n}$ and the left-hand side of (A.4) may be rewritten as follows

$$\begin{aligned} & \mathcal{Q}^{-1} \left[-\frac{d}{dz} \left(\frac{1}{r^2} \mathcal{H} \mathcal{F} \mathcal{H} - \mathcal{F} \right) + [T_4, \mathcal{Q}] - (\mathcal{H} + \mathcal{F}) \left(\frac{1}{r^2} \mathcal{H} \mathcal{F} \mathcal{H} - \mathcal{F} \right) - \left(\frac{1}{r^2} \mathcal{H} \mathcal{F} \mathcal{H} - \mathcal{F} \right) (\mathcal{H} + \mathcal{F}) \right] \mathcal{Q}^{-1} \\ &= \mathcal{Q}^{-1} \left[-\frac{1}{r^2} \mathcal{H} \left(\frac{d\mathcal{F}}{dz} - [T_4, \mathcal{F}] \right) \mathcal{H} + \frac{d\mathcal{F}}{dz} - [T_4, \mathcal{F}] - \frac{1}{r^2} (\mathcal{H} + \mathcal{F}) \mathcal{H} \mathcal{F} \mathcal{H} - \frac{1}{r^2} \mathcal{H} \mathcal{F} \mathcal{H} (\mathcal{H} + \mathcal{F}) \right. \\ & \quad \left. + (\mathcal{H} + \mathcal{F}) \mathcal{F} + \mathcal{F} (\mathcal{H} + \mathcal{F}) \right] \mathcal{Q}^{-1} \\ &= \mathcal{Q}^{-1} \left[\frac{1}{r^2} \mathcal{H} \left(\mathcal{F}^2 + 1_2 \otimes \sum_{i=1}^3 T_i T_i \right) \mathcal{H} - \left(\mathcal{F}^2 + 1_2 \otimes \sum_{i=1}^3 T_i T_i \right) \right. \\ & \quad \left. - \frac{1}{r^2} \mathcal{H}^2 \mathcal{F} \mathcal{H} - \frac{1}{r^2} \mathcal{F} \mathcal{H} \mathcal{F} \mathcal{H} + \mathcal{H} \mathcal{F} + \mathcal{F}^2 - \frac{1}{r^2} \mathcal{H} \mathcal{F} \mathcal{H}^2 - \frac{1}{r^2} \mathcal{H} \mathcal{F} \mathcal{H} \mathcal{F} + \mathcal{F} \mathcal{H} + \mathcal{F}^2 \right] \mathcal{Q}^{-1} \end{aligned}$$

Upon noting that

$$\begin{aligned} \mathcal{Q}^2 &= \left(\frac{1}{r^2} \mathcal{H} \mathcal{F} \mathcal{H} - \mathcal{F} \right)^2 = \frac{1}{r^4} \mathcal{H} \mathcal{F} \mathcal{H}^2 \mathcal{F} \mathcal{H} + \mathcal{F}^2 - \frac{1}{r^2} \mathcal{H} \mathcal{F} \mathcal{H} \mathcal{F} - \frac{1}{r^2} \mathcal{F} \mathcal{H} \mathcal{F} \mathcal{H} \\ &= \frac{1}{r^2} \mathcal{H} \mathcal{F}^2 \mathcal{H} - \frac{1}{r^2} \mathcal{H} \mathcal{F} \mathcal{H} \mathcal{F} - \frac{1}{r^2} \mathcal{F} \mathcal{H} \mathcal{F} \mathcal{H} + \mathcal{F}^2 \end{aligned}$$

and performing the appropriate cancellations we obtain the necessary result. \square

Proposition A.2. *With \mathcal{Q} as previously given we have the antiderivative*

$$(A.5) \quad \int dz z \mathbf{v}_a^\dagger \mathbf{v}_b = \mathbf{v}_a^\dagger \mathcal{S} \mathbf{v}_b, \quad \text{where} \quad \mathcal{S} = \mathcal{Q}^{-1} \left(z + \mathcal{H} \frac{x_i}{r^2} \frac{\partial}{\partial x_i} \right).$$

Proof. Set $\mathcal{S}_1 = \mathcal{Q}^{-1} z$ and $\mathcal{S}_2 = \mathcal{Q}^{-1} \mathcal{H} \frac{x_i}{r^2} \frac{\partial}{\partial x_i}$. Then

$$\mathcal{D}(\mathcal{S}_1) = \frac{d}{dz} (z \mathcal{Q}^{-1}) - z [T_4, \mathcal{Q}^{-1}] - z (\mathcal{H} + \mathcal{F}) \mathcal{Q}^{-1} - z \mathcal{Q}^{-1} (\mathcal{H} + \mathcal{F}) = z \mathcal{D}(\mathcal{Q}^{-1}) + \mathcal{Q}^{-1}$$

$$= z1_{2n} + \mathcal{Q}^{-1}.$$

Further, again using (A.4),

$$\begin{aligned} r^2 \mathcal{D}(\mathcal{S}_2) &= \frac{d}{dz} \left(\mathcal{Q}^{-1} \mathcal{H} x_i \frac{\partial}{\partial x_i} \right) - [T_4, \mathcal{Q}^{-1} \mathcal{H} x_i \frac{\partial}{\partial x_i}] - (\mathcal{H} + \mathcal{F}) \mathcal{Q}^{-1} \mathcal{H} x_i \frac{\partial}{\partial x_i} - \mathcal{Q}^{-1} \mathcal{H} x_i \frac{\partial}{\partial x_i} (\mathcal{H} + \mathcal{F}) \\ &= \mathcal{D}(\mathcal{Q}^{-1}) \mathcal{H} x_i \frac{\partial}{\partial x_i} + \mathcal{Q}^{-1} (\mathcal{H} + \mathcal{F}) \mathcal{H} x_i \frac{\partial}{\partial x_i} - \mathcal{Q}^{-1} \mathcal{H} x_i \frac{\partial}{\partial x_i} (\mathcal{H} + \mathcal{F}) \\ &= \mathcal{H} x_i \frac{\partial}{\partial x_i} + \mathcal{Q}^{-1} (\mathcal{H} + \mathcal{F}) \mathcal{H} x_i \frac{\partial}{\partial x_i} - \mathcal{Q}^{-1} \mathcal{H} (\mathcal{H} + \mathcal{F}) x_i \frac{\partial}{\partial x_i} - \mathcal{Q}^{-1} \mathcal{H} x_i (\partial_i \mathcal{H}) \\ &= \mathcal{Q}^{-1} [\mathcal{Q} \mathcal{H} + (\mathcal{H} + \mathcal{F}) \mathcal{H} - \mathcal{H} (\mathcal{H} + \mathcal{F})] x_i \frac{\partial}{\partial x_i} - \mathcal{Q}^{-1} \mathcal{H} x_i (\partial_i \mathcal{H}) \\ &= -\mathcal{Q}^{-1} r^2 \end{aligned}$$

Upon combining these, $\mathcal{D}(\mathcal{S}) = \mathcal{D}(\mathcal{S}_1 + \mathcal{S}_2) = z1_{2n}$ and the result follows. \square

We note that we may write \mathcal{S}_2) more symmetrically if needed:

$$\mathcal{S}_2 = \frac{1}{2} \left[\overleftarrow{\frac{\partial}{\partial x_i}} \frac{x_i}{r^2} \mathcal{H} \mathcal{Q}^{-1} + \mathcal{Q}^{-1} \mathcal{H} \frac{x_i}{r^2} \overrightarrow{\frac{\partial}{\partial x_i}} \right].$$

Remark: In [15] Panagopolous gave the ordinary differential operator $\mathcal{S}' = \mathcal{Q}^{-1} (z + 2\mathcal{H} \frac{d}{dr^2})$ for which $\mathcal{D}(\mathcal{S}') = z1_{2n} + \mathcal{Q}^{-1} [\frac{d\mathcal{H}}{dr^2}, \mathcal{H}]$; as we are acting on functions and gauge transformations that are not just a function of r we have replaced this with our partial differential operator \mathcal{S} .

Proposition A.3. *With \mathcal{Q} as previously given we have the antiderivative*

$$(A.6) \quad \int \mathbf{v}_a^\dagger \frac{\partial}{\partial x_i} \mathbf{v}_b dz = \mathbf{v}_a^\dagger \mathcal{Q}^{-1} \left[\frac{\partial}{\partial x_i} + \mathcal{H} \frac{z}{r^2} x_i + \mathcal{H} \frac{\mathbf{l}}{r^2} (\mathbf{x} \times \nabla)_i \right] \mathbf{v}_b.$$

Proof. Let $L = L_1 + L_2 + L_3$ with

$$L_1 = \mathcal{Q}^{-1} \frac{\partial}{\partial x_i}, \quad L_2 = \mathcal{Q}^{-1} \mathcal{H} \frac{z}{r^2} x_i, \quad L_3 = \mathcal{Q}^{-1} \mathcal{H} \frac{\mathbf{l}}{r^2} (\mathbf{x} \times \nabla)_i.$$

We compute $\mathcal{D}(L_i)$, $i = 1, 2, 3$. First

$$\begin{aligned} \mathcal{D}(L_1) &= \frac{d}{dz} \left(\mathcal{Q}^{-1} \frac{\partial}{\partial x_i} \right) - [T_4, \mathcal{Q}^{-1} \frac{\partial}{\partial x_i}] - (\mathcal{H} + \mathcal{F}) \mathcal{Q}^{-1} \frac{\partial}{\partial x_i} - \mathcal{Q}^{-1} \frac{\partial}{\partial x_i} (\mathcal{H} + \mathcal{F}) \\ &= \mathcal{D}(\mathcal{Q}^{-1}) \frac{\partial}{\partial x_i} - \mathcal{Q}^{-1} \frac{\partial \mathcal{H}}{\partial x_i} = 1_{2n} \frac{\partial}{\partial x_i} - \mathcal{Q}^{-1} \frac{\partial \mathcal{H}}{\partial x_i} \end{aligned}$$

where we use Proposition A.1. Next

$$\begin{aligned} \mathcal{D}(L_2) &= \frac{d}{dz} \left(\mathcal{Q}^{-1} \mathcal{H} \frac{z}{r^2} x_i \right) - [T_4, \mathcal{Q}^{-1} \mathcal{H} \frac{z}{r^2} x_i] - (\mathcal{H} + \mathcal{F}) \mathcal{Q}^{-1} \mathcal{H} \frac{z}{r^2} x_i - \mathcal{Q}^{-1} \mathcal{H} \frac{z}{r^2} x_i (\mathcal{H} + \mathcal{F}) \\ &= \mathcal{D}(\mathcal{Q}^{-1}) \mathcal{H} \frac{z}{r^2} x_i + \mathcal{Q}^{-1} (\mathcal{H} + \mathcal{F}) \mathcal{H} \frac{z}{r^2} x_i - \mathcal{Q}^{-1} \mathcal{H} \frac{z}{r^2} x_i (\mathcal{H} + \mathcal{F}) + \mathcal{Q}^{-1} \mathcal{H} \frac{x_i}{r^2} \\ &= \mathcal{H} \frac{z}{r^2} x_i + \mathcal{Q}^{-1} [\mathcal{F}, \mathcal{H}] \frac{z}{r^2} x_i + \mathcal{Q}^{-1} \mathcal{H} \frac{x_i}{r^2} \\ &= \mathcal{Q}^{-1} \mathcal{H} \frac{x_i}{r^2} \end{aligned}$$

upon using (A.2). Next we calculate $\mathcal{D}(L_3)$ for $i = 1$. We have

$$\begin{aligned}
\mathcal{D}(L_3) &= \frac{d}{dz} \left[\mathcal{Q}^{-1} \mathcal{H} \frac{\imath}{r^2} \left(x_2 \frac{\partial}{\partial x_3} - x_3 \frac{\partial}{\partial x_2} \right) \right] - \left[T_4, \mathcal{Q}^{-1} \mathcal{H} \frac{\imath}{r^2} \left(x_2 \frac{\partial}{\partial x_3} - x_3 \frac{\partial}{\partial x_2} \right) \right] \\
&\quad - (\mathcal{H} + \mathcal{F}) \left[\mathcal{Q}^{-1} \mathcal{H} \frac{\imath}{r^2} \left(x_2 \frac{\partial}{\partial x_3} - x_3 \frac{\partial}{\partial x_2} \right) \right] - \left[\mathcal{Q}^{-1} \mathcal{H} \frac{\imath}{r^2} \left(x_2 \frac{\partial}{\partial x_3} - x_3 \frac{\partial}{\partial x_2} \right) \right] (\mathcal{H} + \mathcal{F}) \\
&= \mathcal{D}(\mathcal{Q}^{-1}) \mathcal{H} \frac{\imath}{r^2} \left(x_2 \frac{\partial}{\partial x_3} - x_3 \frac{\partial}{\partial x_2} \right) \\
&\quad + \mathcal{Q}^{-1} (\mathcal{H} + \mathcal{F}) \mathcal{H} \frac{\imath}{r^2} \left(x_2 \frac{\partial}{\partial x_3} - x_3 \frac{\partial}{\partial x_2} \right) - \mathcal{Q}^{-1} \mathcal{H} \frac{\imath}{r^2} \left(x_2 \frac{\partial}{\partial x_3} - x_3 \frac{\partial}{\partial x_2} \right) (\mathcal{H} + \mathcal{F}) \\
&= \mathcal{H} \frac{\imath}{r^2} \left(x_2 \frac{\partial}{\partial x_3} - x_3 \frac{\partial}{\partial x_2} \right) + \mathcal{Q}^{-1} [\mathcal{F}, \mathcal{H}] \frac{\imath}{r^2} \left(x_2 \frac{\partial}{\partial x_3} - x_3 \frac{\partial}{\partial x_2} \right) \\
&\quad - \mathcal{Q}^{-1} \mathcal{H} \frac{\imath}{r^2} \left(x_2 \frac{\partial}{\partial x_3} - x_3 \frac{\partial}{\partial x_2} \right) \mathcal{H} \\
&= -\mathcal{Q}^{-1} \mathcal{H} \frac{\imath}{r^2} \left(x_2 \frac{\partial}{\partial x_3} - x_3 \frac{\partial}{\partial x_2} \right) \mathcal{H}.
\end{aligned}$$

again using (A.2). Altogether we have

$$\mathcal{D}(L) = 1_{2n} \frac{\partial}{\partial x_1} + \mathcal{Q}^{-1} \left\{ -\frac{\partial \mathcal{H}}{\partial x_1} + \mathcal{H} \frac{x_1}{r^2} - \mathcal{H} \frac{\imath}{r^2} \left(x_2 \frac{\partial}{\partial x_3} - x_3 \frac{\partial}{\partial x_2} \right) \mathcal{H} \right\}$$

Multiplying the expression in the parentheses by $-r^2$ gives

$$\begin{aligned}
-\{\cdot\} r^2 &= -\sigma_1 \otimes 1_2(x_1^2 + x_2^2 + x_3^2) + \sigma_1 \otimes 1_2 x_1^2 + \sigma_2 \otimes 1_2 x_1 x_2 + \sigma_3 \otimes 1_2 x_1 x_3 \\
&\quad - \imath(\sigma_1 \otimes 1_2 x_1 + \sigma_2 \otimes 1_2 x_2 + \sigma_3 \otimes 1_2 x_3) \times (x_2 \sigma_3 \otimes 1_2 - x_3 \sigma_2 \otimes 1_2)
\end{aligned}$$

which vanishes by standard relations, proving the result. \square

APPENDIX B. THE NAHM ANSATZ

Introduce the shorthand

$$(B.1) \quad R_j = T_j + \imath x_j 1_n, \quad j = 1, 2, 3, \quad \gamma = -\imath[(x_1 - \imath x_2)\zeta - \imath x_3].$$

Upon substituting (3.11) into $\Delta \mathbf{w} = 0$ we find

$$\begin{aligned}
0 &= |s\rangle \otimes \left(\imath \frac{d}{dz} - \imath T_4 + \imath \gamma + \hat{\mathbf{u}} \cdot \mathbf{R} \right) \hat{\mathbf{w}}(z) \\
&\quad + \sigma_k |s\rangle \otimes \left(\imath \hat{u}^k \left(\frac{d}{dz} - T_4 + \gamma \right) + R^k + \imath(\mathbf{R} \times \hat{\mathbf{u}})^k \right) \hat{\mathbf{w}}(z).
\end{aligned}$$

and so we require

$$(B.2) \quad 0 = \left(\imath \frac{d}{dz} - \imath T_4 + \imath \gamma + \hat{\mathbf{u}} \cdot \mathbf{R} \right) \hat{\mathbf{w}}(z),$$

$$(B.3) \quad 0 = \mathcal{L}_k \hat{\mathbf{w}}(z) := \left(\imath \hat{u}^k \left(\frac{d}{dz} - T_4 + \gamma \right) + R^k + \imath(\mathbf{R} \times \hat{\mathbf{u}})^k \right) \hat{\mathbf{w}}(z).$$

The consistency of these equations imposes various constraints. First consider

$$[\mathcal{L}_1, \mathcal{L}_2] = (\imath \hat{u}^1 + \hat{u}^2 \hat{u}^3) \left(\dot{T}_2 - [T_4, T_1] - [T_3, T_1] \right) - (\imath \hat{u}^2 - \hat{u}^1 \hat{u}^3) \left(\dot{T}_1 - [T_4, T_2] - [T_2, T_3] \right)$$

$$-(1 - (\hat{u}^3)^2) \left(\dot{T}_3 - [T_4, T_3] - [T_1, T_2] \right) + (1 - \hat{\mathbf{u}} \cdot \hat{\mathbf{u}}) \left(\dot{T}_3 - [T_4, T_3] \right).$$

Thus provided $\hat{\mathbf{u}}(\mathbf{x})$ is a unit vector and the T_i 's satisfy the Nahm equations we have consistency of the equations $\mathcal{L}_k \hat{\mathbf{w}}(\mathbf{z}) = 0$.

At this stage we introduce a convenient parametrization (reflected in Hitchin's minitwistor construction). Let $\mathbf{y} \in \mathbb{C}^3$ be a null vector. We may consider $\mathbf{y} \in \mathbb{P}^2$ and parameterize \mathbf{y} as

$$(B.4) \quad \mathbf{y} = \left(\frac{1 + \zeta^2}{2\imath}, \frac{1 - \zeta^2}{2}, -\zeta \right).$$

Then

$$\mathbf{y} \cdot \bar{\mathbf{y}} = \frac{(1 + |\zeta|^2)^2}{2}, \quad \mathbf{y} \cdot \mathbf{y} = 0.$$

The signs here have been chosen so that

$$L(\zeta) := 2\imath \mathbf{y} \cdot \mathbf{T} = (T_1 + \imath T_2) - 2\imath T_3 \zeta + (T_1 - \imath T_2) \zeta^2.$$

In due course we will see this to be our Lax matrix. Set

$$(B.5) \quad \hat{\mathbf{u}} = \hat{\mathbf{u}}(\zeta) := \imath \frac{\mathbf{y} \times \bar{\mathbf{y}}}{\mathbf{y} \cdot \bar{\mathbf{y}}} = \frac{1}{1 + |\zeta|^2} (\imath(\zeta - \bar{\zeta}), (\zeta + \bar{\zeta}), 1 - |\zeta|^2).$$

Then

$$\hat{\mathbf{u}} \times \mathbf{y} = -\imath \mathbf{y}, \quad \hat{\mathbf{u}} \times \bar{\mathbf{y}} = \imath \bar{\mathbf{y}}.$$

The three vectors $\text{Re}(\mathbf{y})$, $\text{Im}(\mathbf{y})$ and $\hat{\mathbf{u}}$ form an orthogonal basis in \mathbb{R}^3 with $|\hat{\mathbf{u}}| = 1$, whence any $\mathbf{v} \in \mathbb{R}^3$ may be written as

$$\mathbf{v} = \hat{\mathbf{u}} (\hat{\mathbf{u}} \cdot \mathbf{v}) + \bar{\mathbf{y}} \left(\frac{\mathbf{y} \cdot \mathbf{v}}{\mathbf{y} \cdot \bar{\mathbf{y}}} \right) + \mathbf{y} \left(\frac{\bar{\mathbf{y}} \cdot \mathbf{v}}{\mathbf{y} \cdot \bar{\mathbf{y}}} \right).$$

In particular,

$$(B.6) \quad \mathbf{v} + \imath \mathbf{v} \times \hat{\mathbf{u}} = \hat{\mathbf{u}} (\hat{\mathbf{u}} \cdot \mathbf{v}) + 2\bar{\mathbf{y}} \left(\frac{\mathbf{y} \cdot \mathbf{v}}{\mathbf{y} \cdot \bar{\mathbf{y}}} \right).$$

We record that

$$\begin{aligned} \overline{\mathbf{y}(\zeta)} &= -\bar{\zeta}^2 \mathbf{y}(-1/\bar{\zeta}), & \hat{\mathbf{u}}(-1/\bar{\zeta}) &= -\hat{\mathbf{u}}(\zeta), \\ \hat{\mathbf{u}} &= (-\imath \zeta^{-1}, \zeta^{-1}, -1) - \frac{2\mathbf{y}}{\zeta(1 + |\zeta|^2)} = (\imath \zeta, \zeta, 1) + \frac{2\bar{\zeta}\bar{\mathbf{y}}}{1 + |\zeta|^2}, \\ \hat{\mathbf{u}} \cdot \mathbf{T} &= -\imath [(T_1 + \imath T_2)\zeta^{-1} - \imath T_3] - \frac{2\mathbf{y} \cdot \mathbf{T}}{\zeta(1 + |\zeta|^2)} = \imath [(T_1 - \imath T_2)\zeta - \imath T_3] + \frac{2\bar{\zeta}\bar{\mathbf{y}} \cdot \mathbf{T}}{1 + |\zeta|^2}. \end{aligned}$$

Parameterizing $\hat{\mathbf{u}}$ as above and using (B.6) we may write

$$\begin{aligned} \imath \hat{\mathbf{u}} \left(\frac{d}{dz} - T_4 + \gamma \right) + \mathbf{R} + \imath \mathbf{R} \times \hat{\mathbf{u}} &= \imath \hat{\mathbf{u}} \left(\frac{d}{dz} - T_4 + \gamma \right) + \hat{\mathbf{u}} (\hat{\mathbf{u}} \cdot \mathbf{R}) + 2\bar{\mathbf{y}} \left(\frac{\mathbf{y} \cdot \mathbf{R}}{\mathbf{y} \cdot \bar{\mathbf{y}}} \right) \\ &= \hat{\mathbf{u}} \left(\imath \left(\frac{d}{dz} - T_4 + \gamma \right) + \hat{\mathbf{u}} \cdot \mathbf{R} \right) + 2\bar{\mathbf{y}} \left(\frac{\mathbf{y} \cdot \mathbf{R}}{\mathbf{y} \cdot \bar{\mathbf{y}}} \right) \end{aligned}$$

and as a consequence (B.2, B.3) are equivalent to

$$(B.7) \quad 0 = \left(\imath \frac{d}{dz} - \imath T_4 + \imath \gamma + \hat{\mathbf{u}} \cdot \mathbf{R} \right) \hat{\mathbf{w}}(z),$$

$$(B.8) \quad 0 = (\mathbf{y} \cdot \mathbf{R}) \hat{\mathbf{w}}(z).$$

The remaining consistency to be checked is then

$$\left[\imath \frac{d}{dz} - \imath T_4 + \imath \gamma + \hat{\mathbf{u}} \cdot \mathbf{R}, \mathbf{y} \cdot \mathbf{R} \right] = \imath \mathbf{y} \cdot \left(\dot{\mathbf{T}} - [T_4, \mathbf{T}] \right) + [\hat{\mathbf{u}} \cdot \mathbf{T}, \mathbf{y} \cdot \mathbf{T}] = 0,$$

which upon use of $\hat{\mathbf{u}} \times \mathbf{y} = -\imath \mathbf{y}$ is equivalent to Nahm's equations.

Equally from

$$\hat{\mathbf{u}} \cdot \mathbf{R} = -\imath [(R_1 + \imath R_2)\zeta^{-1} - \imath R_3] - \frac{2\mathbf{y} \cdot \mathbf{R}}{\zeta(1 + |\zeta|^2)} = \imath [(R_1 - \imath R_2)\zeta - \imath R_3] + \frac{2\bar{\zeta}\mathbf{y} \cdot \mathbf{R}}{1 + |\zeta|^2}$$

we may write the equations as

$$\begin{aligned} 0 &= \left(\frac{d}{dz} - T_4 + \gamma + [(R_1 - \imath R_2)\zeta - \imath R_3] \right) \hat{\mathbf{w}}(z) = \left(\frac{d}{dz} + M \right) \hat{\mathbf{w}}(z), \\ 0 &= (\mathbf{y} \cdot \mathbf{R}) \hat{\mathbf{w}}(z), \end{aligned}$$

where

$$(B.9) \quad M(\zeta) := (T_1 - \imath T_2)\zeta - \imath T_3 - T_4.$$

The equations we have obtained are just the Lax equations

$$\begin{aligned} 0 &= 2\imath (\mathbf{y} \cdot \mathbf{R}) \hat{\mathbf{w}}(z) = (L(\zeta) - \eta) \hat{\mathbf{w}}(z), \\ 0 &= \left(\frac{d}{dz} + M \right) \hat{\mathbf{w}}(z), \end{aligned}$$

for which $\dot{L} = [L, M]$ and where our construction defines η to be

$$\eta = 2\mathbf{y} \cdot \mathbf{x} = (x_2 - \imath x_1) - 2x_3\zeta - (x_2 + \imath x_1)\zeta^2.$$

From the first of these we see that

$$0 = \det(L(\zeta) - \eta),$$

which gives the equation of the spectral curve \mathcal{C} . Upon using $\overline{\mathbf{y}(\zeta)} = -\bar{\zeta}^2 \mathbf{y}(-1/\bar{\zeta})$ we see from

$$0 = \det(L(\zeta) - \eta)^\dagger = \det(L(\zeta)^\dagger - \bar{\eta}) = \det(2\imath \overline{\mathbf{y}(\zeta)} \cdot \mathbf{T} - \bar{\eta}) = \det(-2\imath \bar{\zeta}^2 \mathbf{y}(-1/\bar{\zeta}) \cdot \mathbf{T} - \bar{\eta})$$

that the spectral curve is invariant under

$$\mathfrak{J} : (\zeta, \eta) \rightarrow \left(-\frac{1}{\bar{\zeta}}, -\frac{\bar{\eta}}{\bar{\zeta}^2} \right).$$

The spectral curve then has the form

$$(B.10) \quad P(\eta, \zeta) := \eta^n + a_1(\zeta)\eta^{n-1} + \dots + a_n(\zeta) = 0, \quad \deg a_k(\zeta) \leq 2k,$$

and the genus of \mathcal{C} is $g = (n-1)^2$.

APPENDIX C. DIRECT INTEGRATION OF THE LAX EQUATIONS

When Nahm gave his ansatz he also suggested a direct integration of the resulting Lax equations; Panagopoulos [15, §3] tried implementing this for the charge 2 case. We will review here this approach and conclude by contrasting it with the proposed Baker-Akhiezer approach we have described.

Underlying the algebro-geometric description of Lax pairs and the spectral curve is that $\dim_{\mathbb{C}} \ker [L(\zeta) - \eta] = 1$. Thus if \mathbf{f}_i is an eigenvector of L with associated eigenvalue η_i then so too is $\mathbf{f}_i h_i(z)$. The idea is that given a solution of (3.12) to find an appropriate $h_i(z)$ such that (3.13) is also satisfied. To implement this we need to know both an eigenvector

\mathbf{f}_i and its derivative to subsequently obtain a differential equation for $h_i(z)$. To determine these we make some simple observations:

- (1) Suppose $L := L(\zeta)$ satisfies the Lax equation $\dot{L} = [L, M]$. Then for any (ζ, η)

$$\det(L(\zeta) - \eta) 1_n = (L(\zeta) - \eta) \text{Adj}(L(\zeta) - \eta)$$

is constant, where Adj denotes the adjugate matrix. Differentiating this yields

$$(L(\zeta) - \eta) \left(\frac{d}{dz} \text{Adj}(L(\zeta) - \eta) - [\text{Adj}(L(\zeta) - \eta), M] \right) = 0$$

and using the generic invertibility of $L(\zeta) - \eta$ we obtain

$$(C.1) \quad \frac{d}{dz} \text{Adj}(L(\zeta) - \eta) = [\text{Adj}(L(\zeta) - \eta), M].$$

- (2) For any constant vector $\boldsymbol{\nu}$ let $P = (\zeta, \eta_i)$ lie on the spectral curve. Then

$$(C.2) \quad \mathbf{f}_i = \text{Adj}(L - \eta_i) \boldsymbol{\nu}$$

is an eigenvector of $L(\zeta)$ with eigenvalue η_i as

$$(L - \lambda_i) \mathbf{f}_i = (L - \eta_i) \text{Adj}(L - \eta_i) \boldsymbol{\nu} = \det(L - \eta_i) \boldsymbol{\nu} = 0.$$

Using these observations we seek a function $h_i(z)$ such that $F = \text{Adj}(L - \eta_i) \boldsymbol{\nu} h_i(z)$ is solution of (3.13). Now

$$\begin{aligned} \dot{F} &= [\text{Adj}(L - \eta_i), M] \boldsymbol{\nu} h_i(z) + \text{Adj}(L - \eta_i) \boldsymbol{\nu} \dot{h}_i(z) \\ &= -MF + F h_i^{-1} \dot{h}_i + \text{Adj}(L - \eta_i) M \boldsymbol{\nu} h_i(z) \end{aligned}$$

Taking the inner product with an arbitrary vector $\boldsymbol{\mu}$ and requiring F satisfy (3.13) yields the differential equation

$$(C.3) \quad h_i^{-1} \frac{dh_i}{dz} = - \frac{\boldsymbol{\mu}^T \text{Adj}(L - \eta_i) M \boldsymbol{\nu}}{\boldsymbol{\mu}^T \text{Adj}(L - \eta_i) \boldsymbol{\nu}}.$$

We see then that to employ this approach we need both the Nahm data to be able to determine the right-hand side of (C.3) and to be able to integrate the subsequent rational expressions. For $n = 2$ this is possible. For higher charges we can only construct Nahm data up to the gauge transformation described in the main text. Let us imagine we make such a gauge transformation and then use the approach of this appendix to solve (3.12, 3.13). Now the right-hand side of (C.3) is in principle known and an integration is required. By way of contrast the Baker-Akhiezer approach allows the construction of the gauge transform of both L and the solution of (3.13) without further integration.

REFERENCES

- [1] H. W. Braden, Antonella D’Avanzo, and V. Z. Enolski. On charge-3 cyclic monopoles. *Nonlinearity*, 24(3):643–675, 2011.
- [2] H. W. Braden and V. Z. Enolski. A Platonic monopole, a related Riemann surface, and an identity of Ramanujan. *Ukrain. Fiz. Zh.*, 50(8):866–874, 2005.
- [3] H. W. Braden and V. Z. Enolski. On the tetrahedrally symmetric monopole. *Comm. Math. Phys.*, 299(1):255–282, 2010.
- [4] H. W. Braden and Yu. N. Fedorov. An extended Abel-Jacobi map. *J. Geom. Phys.*, 58(10):1346–1354, 2008.
- [5] W. Braden and V. Z. Enolski. Some remarks on the Ercolani-Sinha construction of monopoles. *Theoretical and Mathematical Physics*, 165(3):5671597, 2010.
- [6] S. K. Donaldson. Nahm’s equations and the classification of monopoles. *Comm. Math. Phys.*, 96(3):387–407, 1984.
- [7] N. Ercolani and A. Sinha. Monopoles and Baker functions. *Comm. Math. Phys.*, 125(3):385–416, 1989.

- [8] Peter Forgács, Zolán Horváth, and László Palla. Finitely separated multimonopoles generated as solitons. *Phys. Lett. B*, 109(3):200–204, 1982.
- [9] N. J. Hitchin. On the construction of monopoles. *Comm. Math. Phys.*, 89(2):145–190, 1983.
- [10] Jacques Hurtubise. SU(2) monopoles of charge 2. *Comm. Math. Phys.*, 92(2):195–202, 1983.
- [11] Jacques Hurtubise. The asymptotic Higgs field of a monopole. *Comm. Math. Phys.*, 97(3):381–389, 1985.
- [12] I. M. Kričever. Integration of nonlinear equations by the methods of algebraic geometry. *Funkcional. Anal. i Prilozhen.*, 11(1):15–31, 96, 1977.
- [13] W. Nahm. On abelian self-dual multimonopoles. *Phys. Lett. B*, 93(1-2):42–46, 1980.
- [14] Werner Nahm. The algebraic geometry of multi-monopoles. In *Group Theoretical Methods in Physics. Proceedings, 11th International Colloquium, Istanbul, Turkey, August 23-28, 1982*, 1982.
- [15] H. Panagopoulos. Multimonopoles in arbitrary gauge groups and the complete SU(2) two-monopole system. *Phys. Rev. D (3)*, 28(2):380–384, 1983.
- [16] R. S. Ward. Ansätze for self-dual Yang-Mills fields. *Comm. Math. Phys.*, 80(4):563–574, 1981.
- [17] Erick J. Weinberg and Piljin Yi. Magnetic monopole dynamics, supersymmetry, and duality. *Phys. Rep.*, 438(2-4):65–236, 2007.

SCHOOL OF MATHEMATICS, EDINBURGH UNIVERSITY, EDINBURGH.

E-mail address: hwb@ed.ac.uk

NATIONAL UNIVERSITY OF “KYIV-MOHYLA ACADEMY”, KYIV, UKRAINE, 04655.

E-mail address: venolski@googlemail.com